

## Institute for Animal Health

Registered Office: Compton, Newbury, Berkshire RG20 7NN Telephone: +44 (0) 1635-578411 Facsimile: +44 (0) 1635-577237

e-mail: iah@bbsrc.ac.uk Website: http://www.iah.bbsrc.ac.uk

Director: Professor Martin Shirley PhD

Reply to Pirbright Laboratory Fax: 01483 232621

Direct Dial: 01483 231014

E-mail: Elizabeth.byrom@bbsrc.ac.uk

Page 1 of 2

To:

Email:

CC:

Fax Number:

From:

30th March 2009 Date:

Subject: R1 Report

No. Of Pages: 2

This message is intended for the use of the person named above and may contain confidential information. Any other distribution, copying or disclosure is strictly prohibited. If you have received this telefax in error, please notify us immediately by telephone or E-mail and then return the original transmission to us by mail.

Thank you.

Dear

Please find below the final "r1" value report for A Iraq 21/2009 and A Iraq 24/2009.

Yours sincerely,

Head: World Reference Laboratory for FMD

nce No. 228824 The Institute is sponsored by the Biotechnology and Biological Sciences Research Council. An Associated Institute of the University of Reading.

Pirbright Laboratory Ash Road, Pirbright, Woking Surrey GU24 0NF Tel: 01483-232441 Fax: 01483-232448

Compton Laboratory Compton, Newbury, Berkshire RG20 7NN Tel: 01635-578411

Fax: 01635-577237

Report no:	07/09	2dmVNT							LPBE			
Field Isolate:	SAU Isolate ref:	2dmVNT test ref:	A Tur06 Arriah	A22 lrq 23-32	A22 lrq 34-43	A Sau 41/91 SI94	A Eri RZ pool	A Ind 17/82 7756	ELISA test ref:	A22 Irq 24/64 TC76-83	A May 97 VJ62-76	A Irn 87 Merial 1781
A Irq 21/2009	B83/09	mean	0.60	0.11	0.07	0.19	poor	7730	mean	0.08	0.83	0.05
A Irq 24/2009	B84/09		0.81	0.53	0.50	0.24	≤0.04	0.22	mean	0.50	≥1	0.15

## Interpretation of r<sub>1</sub> values

## In the case of ELISA:

 $r_1 = 0.4-1.0$ . Suggests that there is a close relationship between field isolate and vaccine strain. A potent vaccine containing the vaccine strain is likely to confer protection.

 $r_1 = 0.2$ -0.39, Suggests that the field isolate is antigenically related to the vaccine strain. The vaccine strain might be suitable for use if no closer match can be found provided that a potent vaccine is used and animals are preferably immunised more than once.

 $r_1 = <0.2$ . Suggests that the field isolate is so different from the vaccine strain that the vaccine is unlikely to protect.

## In the case of neutralisation:

 $r_1 = \ge 0.3$ . Suggests that there is a close relationship between field isolate and vaccine strain. A potent vaccine containing the vaccine strain is likely to confer protection.

 $r_1 = < 0.3$ . Suggests that the field isolate is so different from the vaccine strain that the vaccine is unlikely to protect.

N.B.

All of our phylogenetic trees can be accessed via the internet at:

http://www.iah.bbsrc.ac.uk/primary\_index/current\_research/virus/Picornaviridae/Aphthovirus/index.html